# HOW TO DISCUSS YOUR ACUTE HEPATIC PORPHYRIA (AHP) WITH YOUR HEALTHCARE PROVIDER

Your answers to this questionnaire will help you better communicate your experience with AHP. Print out this guide and *bring it to your next doctor's appointment* to start a conversation about your history with AHP and how it currently impacts your life.

1.	What t	ype of A	AHP do	you h	nave?

Acute intermittent porphyria (AIP)

Hereditary coproporphyria (HCP)

Variegate porphyria (VP)

ALAD-deficiency porphyria (ADP)

2.	When were	you diagnosed	with AHP?

Month: Yea	ar:
------------	-----

3.	Please rate the average day:	•	of your	acute hepa	tic porphy	ria (AHP) syr	nptoms on an
	1	2	3	4	5		
	Least Severe			N	Nost Severe		
4.	What sympto	oms of AHP	do you	experience	eș.		
5.	In the past m		nany da	ays have you	ur sympto	ms of AHP fe	lt worse
	0-5 days	6-10 days	5	11-15 days	16	+ days	
6.	In the past m		nany da	ays have you	ur sympto	ms of AHP fe	lt better
	0-5 days	6-10 days	5	11-15 days	16	+ days	
7.	How many ti urgent healtl	•	-	•	•		
	0 1	2	3	4	5+		
8.	How does Al	HP disrupt y	our dai	ly life and a	ctivities?		
9.	Which of the	ese disrupte	d activi	ties affect y	our life th	e most?	

-					
What	do you hope th	nat treatment	can help w	ith the most?	
-	/				
Write	down any addi	tional questic	ons or infor	mation for yo	ur appointment.

Bring this guide with you to your next doctor appointment to discuss your acute hepatic porphyria (AHP) and to see if GIVLAARI® (givosiran) may be a treatment option.

For more information, visit <u>www.GIVLAARI.com</u>.

## What is GIVLAARI® (givosiran)?

GIVLAARI is a prescription medicine used to treat acute hepatic porphyria (AHP) in adults.

# **Important Safety Information**

Do not use GIVLAARI if you have ever had a severe allergic reaction to GIVLAARI.

Please see full Important Safety Information on page 4 and full Prescribing Information.

This discussion guide is not validated by any medical organization and does not replace the opinion of a trained medical physician.

# IMPORTANT SAFETY INFORMATION

Do not use GIVLAARI® (givosiran) if you have ever had a severe allergic reaction to GIVLAARI.

#### **GIVLAARI** can cause:

#### Severe allergic reaction

Tell your doctor or nurse right away if you experience any of the following signs or symptoms of a severe allergic reaction during treatment:

- Swelling mainly of the lips, tongue or throat which makes it difficult to swallow or breathe
- Breathing problems or wheezing
- Rash or hives
- Itching
- · Feeling dizzy or fainting

If you have a severe allergic reaction, your doctor or nurse will stop GIVLAARI treatment right away and you may need to take other medicines to control the symptoms.

#### Liver problems

Your doctor will check your liver function by doing blood tests:

 Before you start using GIVLAARI

- Once a month for the first
  6 months of treatment
- And when they think it is needed

If these tests show abnormal results, your doctor or nurse will decide whether to temporarily interrupt or stop treatment with GIVLAARI.

#### Kidney problems

Your doctor will check how your kidneys are working while you are using GIVLAARI.

### • Injection site reactions

GIVLAARI is given as an injection under the skin (called a "subcutaneous injection"). Reactions to this injection may happen during treatment with GIVLAARI.

Tell your doctor or nurse right away if you experience any of the following symptoms of an injection site reaction during treatment: redness, pain, itchiness, rash, discoloration, or swelling around the injection site.

# • Increased blood homocysteine levels

GIVLAARI may cause increased levels of homocysteine (a type of amino acid) in your blood. Your doctor will check your homocysteine levels before and during treatment by doing blood tests. If your levels are increased, your doctor may check your folate, vitamins B12 and B6, and tell you to take a vitamin B6 supplement.

#### What are the common side effects of GIVLAARI?

The most common side effects of GIVLAARI are nausea and injection site reactions. These are not all the possible side effects of GIVLAARI. Talk to your doctor about side effects that you experience. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

For additional information about GIVLAARI, please see full Prescribing Information.



